



## Turf Services Performed Today

\_\_\_\_/\_\_\_\_/\_\_\_\_

### Round #2: Please Stay Off Treated Areas Until Dry

\_\_\_\_\_ **Fertilization w/ Pre-Emergent:** A balanced granular product containing fast and slow release nutrients combined with a pre-emergent herbicide for weed control was applied to your turf. **This application NEEDS TO BE watered within 1-2 weeks for best results.** About ½ inch of rainfall or water is recommended.

\_\_\_\_\_ **Broadleaf Weed Management:** A liquid selective herbicide was applied to broadleaf weeds. This product will usually be effective upon drying within 1-3 hours. **It is recommended that you DO NOT water immediately after application for best results.**

#### **Beneficial Applications If Performed:**

\_\_\_\_\_ **Core Aeration:** Breaks through thatch relieving compaction and opens up soil to allow for air, water, and nutrients get where needed the most. Greatly enhances seed to soil contact and root development.

\_\_\_\_\_ **Disease Management:** A granular or liquid fungicide was applied to some or all of your turf areas due to evidence of or probable development of turf disease. This product will usually be effective upon drying. **It is recommended that you DO NOT water immediately after application for best results.**

#### **Round #2 Turf Care Notes:**

The fertilizer/pre-emergent combination applied will provide nutrients for your lawn and will prevent most weeds, including crabgrass, from becoming established. The herbicide spray is typically an over-spray to control common broadleaf weeds that are present. Insect and disease problems are not usually serious issues in spring. Disease such as “Red Thread” and “Dollar Spot” may appear in late April through June in our area. These may develop after periods of dampness and cool temperatures followed by sunny days but are not typically problematic long-term.

Spring rains, on average, should supply all the watering needs for your lawn. **If you irrigate/water no more than once a week to provide a quarter to a half inch of water per week.** Set irrigation schedules for 1-2 days after mowing. If your lawn appears purplish or bluish-gray and grass blades do not spring back quickly after any type of traffic, your grass is beginning the first phase of drought stress. An immediate deep watering of these areas should bring them back to a healthy condition within hours to 1-2 days.

**If disease is indicated below**, allow turf to completely dry out for 1-2 weeks and cutback on the frequency of watering.

**If stress is indicated below**, please water the areas as deeply as possible each day or so until they recover.

\_\_\_ **Disease Present:** \_\_\_ Red Thread \_\_\_ Dollar Spot \_\_\_ Brown Patch \_\_\_ Other

\_\_\_ **Turf Drought Stress:** \_\_\_ Front \_\_\_ Back \_\_\_ Left Side \_\_\_ Right Side

**Please contact Client Services if you have any questions regarding this service.**

**THANK YOU**

20098 Ashbrook Place / Suite 185 / Ashburn, VA 20147

Call: 703-849-0713 / Fax: 703-849-0728

Write: [customerservice@khlandscaping.com](mailto:customerservice@khlandscaping.com)

Visit: [khlandscaping.com](http://khlandscaping.com)